

Understanding light quanta: First quantization of the free electromagnetic field

A. C. de la Torre

Departamento de Física, Universidad Nacional de Mar del Plata

Funes 3350, 7600 Mar del Plata, Argentina

dltorre@mdp.edu.ar

Abstract

The quantization of the electromagnetic field in vacuum is presented without reference to lagrangean quantum field theory. The equal time commutators of the fields are calculated from basic principles. A physical discussion of the commutators suggest that the electromagnetic fields are macroscopic emergent properties of more fundamental physical system: the photons.

I. INTRODUCTION

In a letter to his friend Besso in 1951, Einstein complained that after fifty years of conscious meditation he had not come any closer to the answer to the question *What are light quanta?* The level of understanding of quantum electrodynamics reached at that time by the physics community was not sufficient for him since he added: “today any brute believes that he knows the answer, but he is wrong”¹. In this work and in the following ones we will present the understanding that we have today on the quantum of electromagnetic radiation. This understanding, perhaps, would still be unsatisfactory for Einstein but at least we can claim that we understand the light quanta as much, or as little, as we understand any other material particle. The main idea in this search for clarity will be to recognize that the electromagnetic fields are nor a primary physical entity but are a collective manifestation of more fundamental entities, the photons, carriers of energy, momentum and angular momentum.

It is remarkable that in almost all quantum mechanics texts, even in the advanced ones, the quantization of the electromagnetic fields is avoided and the subject is postponed to lagrangean quantum field theory. In few cases² an introduction to quantum field theory is given just in order to present the (second) quantization of the electromagnetic field. After having learned about the success and power of quantum mechanics in the description

of physical systems, one should wonder whether the same procedures can be successfully applied to the electromagnetic fields. A didactic relevance of this work is to show that it is possible, and easy, to quantize the free electromagnetic field without any reference to quantum field theory which is, indeed, a difficult subject. However, after seeing that the quantization of the electromagnetic field can, in principle, be performed following the same line of thoughts as, say, the harmonic oscillator, we will give arguments indicating that this is not a meaningful thing to do because there are good reasons for thinking that the electromagnetic field is not a primary physical system, that necessarily requires a quantum treatment, but should be understood as a collective manifestation of an ensemble of more fundamental quantum mechanical objects: the photons.

In this work we will consider only observable quantities like the electric and magnetic fields and we will avoid the use of unobservable potential fields that present considerable complications in a Lorentz invariant gauge fixing adequate for quantization.

II. THE CLASSICAL FREE ELECTROMAGNETIC FIELD

For the description of the main features of the classical free electromagnetic field we adopt Heaviside system, that is, rationalized Gauß units³. The six observables of the physical system are the electric field vector $\mathbf{E}(\mathbf{r}, t)$ and the magnetic field (pseudo)vector $\mathbf{B}(\mathbf{r}, t)$ in vacuum. These fields satisfy Maxwell's equations

$$-\varepsilon_{jkl}\partial_k E_l = \frac{1}{c}\partial_t B_j, \quad (1)$$

$$\varepsilon_{jkl}\partial_k B_l = \frac{1}{c}\partial_t E_j, \quad (2)$$

$$\partial_k E_k = 0, \quad (3)$$

$$\partial_k B_k = 0, \quad (4)$$

where ε_{jkl} is the well known total antisymmetric (pseudo)tensor, $\partial_k = \frac{\partial}{\partial x_k}$, $\partial_t = \frac{\partial}{\partial t}$ and we adopt Einstein's convention, stating that repeated indices indicate a summation. Furthermore, in order to keep the notation as simple as possible we will suppress the space-time arguments of the fields wherever they are not explicitly needed. At any point in space where the fields don't vanish there is an energy and momentum *density* given by the local value of the fields; the *total* energy H and momentum $\mathbf{P}$ are given by the volume integrals

$$H = \frac{1}{8\pi} \int d^3\mathbf{r} (\mathbf{E}^2 + \mathbf{B}^2) \quad (5)$$

$$\mathbf{P} = \frac{1}{8\pi c} \int d^3\mathbf{r} (\mathbf{E} \times \mathbf{B} - \mathbf{B} \times \mathbf{E}). \quad (6)$$

Note that, although at the classical level it is unnecessary, in the definition of momentum $\mathbf{P}$ we allow for a possible noncommutation of the fields. In similar manner, we could define the centroid of the energy distribution and the angular momentum carried by the fields.

In our treatment of electromagnetism we will make intensive use of the transformations of space inversion $\mathcal{P}$, time inversion $\mathcal{T}$, charge conjugation $\mathcal{C}$ and the *dual* transformation $\mathcal{D}$ (defined below). The fields have following transformation properties.

$$\mathcal{P} [\mathbf{E}(\mathbf{r}, t)] = -\mathbf{E}(-\mathbf{r}, t) , \quad (7)$$

$$\mathcal{T} [\mathbf{E}(\mathbf{r}, t)] = \mathbf{E}(\mathbf{r}, -t) , \quad (8)$$

$$\mathcal{C} [\mathbf{E}(\mathbf{r}, t)] = -\mathbf{E}(\mathbf{r}, t) , \quad (9)$$

$$\mathcal{D} [\mathbf{E}(\mathbf{r}, t)] = -\mathbf{B}(\mathbf{r}, t) , \quad (10)$$

$$\mathcal{P} [\mathbf{B}(\mathbf{r}, t)] = \mathbf{B}(-\mathbf{r}, t) , \quad (11)$$

$$\mathcal{T} [\mathbf{B}(\mathbf{r}, t)] = -\mathbf{B}(\mathbf{r}, -t) , \quad (12)$$

$$\mathcal{C} [\mathbf{B}(\mathbf{r}, t)] = -\mathbf{B}(\mathbf{r}, t) , \quad (13)$$

$$\mathcal{D} [\mathbf{B}(\mathbf{r}, t)] = \mathbf{E}(\mathbf{r}, t) . \quad (14)$$

Since the set of (vacuum) Maxwell's equations is invariant under these transformations, the transformation of any true relation among the fields results in another true relation.

It is very convenient to introduce a complex electromagnetic field $\mathbf{F}(\mathbf{r}, t)$ defined by

$$\mathbf{F}(\mathbf{r}, t) = \mathbf{E}(\mathbf{r}, t) + i\mathbf{B}(\mathbf{r}, t) . \quad (15)$$

It has been claimed⁴ that this is not only a very convenient notation but it is a manifestation of a deep mathematical structure, called Geometric Algebra, adequate for modelling physical reality. Note that the real part of $\mathbf{F}$ is a vector and the imaginary part is a pseudovector and that the relation above can be trivially inverted in order to recover the electric and magnetic field that may be more familiar to the reader. For the components of the free electromagnetic field F_k , Maxwell's equations are

$$\varepsilon_{jkl} \partial_k F_l = i \frac{1}{c} \partial_t F_j , \quad (16)$$

$$\partial_k F_k = 0 , \quad (17)$$

and the transformation properties are

$$\mathcal{P} [\mathbf{F}(\mathbf{r}, t)] = -\mathbf{F}^*(-\mathbf{r}, t) , \quad (18)$$

$$\mathcal{T} [\mathbf{F}(\mathbf{r}, t)] = \mathbf{F}^*(\mathbf{r}, -t) \text{ (classical)} , \quad (19)$$

$$\mathcal{C} [\mathbf{F}(\mathbf{r}, t)] = -\mathbf{F}(\mathbf{r}, t) , \quad (20)$$

$$\mathcal{D} [\mathbf{F}(\mathbf{r}, t)] = i\mathbf{F}(\mathbf{r}, t) . \quad (21)$$

The reason for emphasizing that the behaviour of $\mathbf{F}$ under time inversion is classic will be clarified later.

III. FROM CLASSICAL OBSERVABLES TO OPERATORS

We can now follow the standard procedure for the quantum mechanical treatment of a physical system, but applied to the free electromagnetic field. The first step is to define a set of hermitian *field operators* $\mathbf{E}(\mathbf{r}, t)$ and $\mathbf{B}(\mathbf{r}, t)$ in a Hilbert space, corresponding to the electric and magnetic field observables (we use the same symbols to denote the operators and the classical variables but this should cause no confusion). In the Heisenberg picture, the infinite set of operators is parameterized by the variables $(\mathbf{r}, t)$ that are not dynamical variables but are coordinates that characterize a space-time point. Therefore at every space-time point $(\mathbf{r}, t)$ we have six operators $\mathbf{E}$ and $\mathbf{B}$ and we must find the commutation relations for this infinite set. The commutation relations should determine the eigenvalue spectrum related to the observations and the eigenvectors related to the corresponding states of the system.

In the derivation of the commutation relations, we will assume that all relations and symmetries of the classical electromagnetic fields are also satisfied by the corresponding quantum operators. Every constraint on the classical fields should also be present in the quantum description of the fields. The opposite is not true. There are restrictions imposed in the quantum treatment of a system that do not appear in a classical modelling. Quantum mechanics has stronger constraints and correlations that have manifestation in characteristic quantum effects like nonseparability, contextuality, uncertainty relations among classically unrelated quantities, etc. Clearly, the imposition of all classical relations and symmetries is not sufficient to determine the commutation relations. If it were so, in sense, quantum mechanics could be derived from classical mechanics; therefore somewhere we must introduce a quantization postulate that in our case will appear as a mathematical *ansatz* concerning the form of the commutators.

In terms of the field $\mathbf{F}$, the hamiltonian and the momentum operators are given by

$$H = \frac{1}{16\pi} \int d^3\mathbf{r} (\mathbf{F}^\dagger \cdot \mathbf{F} + \mathbf{F} \cdot \mathbf{F}^\dagger) , \quad (22)$$

$$\mathbf{P} = \frac{-i}{16\pi c} \int d^3\mathbf{r} (\mathbf{F}^\dagger \times \mathbf{F} - \mathbf{F} \times \mathbf{F}^\dagger) , \quad (23)$$

where $\mathbf{F}^\dagger = \mathbf{E} - i\mathbf{B}$ is the hermitian adjoint of $\mathbf{F}$ and the noncommutation of the fields has been taken in to account. Using the index notation we have

$$H = \frac{\delta_{kl}}{16\pi} \int d^3\mathbf{r} (F_k^\dagger F_l + F_l F_k^\dagger) , \quad (24)$$

$$P_j = \frac{-i\varepsilon_{jkl}}{16\pi c} \int d^3\mathbf{r} (F_k^\dagger F_l + F_l F_k^\dagger) . \quad (25)$$

The same as is the case in classical physics, the hamiltonian and the momentum are the *generators* of translations in time and space. This means that the space and time derivatives of a field operator (an operator depending on $(\mathbf{r}, t)$) are given by the commutator with momentum and with the hamiltonian respectively⁵. That is

$$\frac{1}{c} \partial_t \circ = \frac{i}{\hbar c} [H, \circ] , \quad (26)$$

$$\partial_j \circ = \frac{1}{i\hbar} [P_j, \circ] . \quad (27)$$

Where the symbol $\circ$ denotes a space holder to be occupied by any field operator. Note that even though these two relations are formally similar, they have completely different origin. The first relation follows from the time evolution of an operator in Heisenberg picture whereas the second is a consequence of the canonical commutation relation between momentum and the position observable. Using these derivatives in Eqs. (16,17), Maxwell's equations for the electromagnetic field operators become

$$(\delta_{jk}\delta_{ls} - \delta_{jl}\delta_{sk} + \delta_{js}\delta_{kl})M_{kls}(\mathbf{r}, t) = 0 , \quad (28)$$

$$\varepsilon_{kls}M_{kls}(\mathbf{r}, t) = 0 , \quad (29)$$

where

$$M_{kls}(\mathbf{r}, t) = \int d^3\mathbf{r}' [F_k^\dagger(\mathbf{r}', t)F_l(\mathbf{r}', t) + F_l(\mathbf{r}', t)F_k^\dagger(\mathbf{r}', t) , F_s(\mathbf{r}, t)] . \quad (30)$$

In the derivation of this result we have used the important identity $\varepsilon_{jkl}\varepsilon_{jsu} = \delta_{ks}\delta_{lu} - \delta_{ku}\delta_{ls}$. We will give a much simpler expression for $M_{kls}(\mathbf{r}, t)$ after obtaining an expression for the fields commutators.

IV. ELECTROMAGNETIC FIELD COMMUTATORS

We will first obtain an expression for the equal-time field commutators derived from a quantization *ansatz* and the requirement of invariance under the $\mathcal{P}, \mathcal{T}, \mathcal{C}$ and $\mathcal{D}$ transformations. Furthermore we will use the condition that physical space is homogeneous and isotropic. Once we know the equal-time commutators we can, in principle, determine the commutators for different times performing a time evolution on one of the fields. The transformation behaviour of the field *operators* are given by

$$\mathcal{P} [\mathbf{F}(\mathbf{r}, t)] = -\mathbf{F}^\dagger(-\mathbf{r}, t) , \quad (31)$$

$$\mathcal{T} [\mathbf{F}(\mathbf{r}, t)] = \mathbf{F}(\mathbf{r}, -t) , \quad (32)$$

$$\mathcal{C} [\mathbf{F}(\mathbf{r}, t)] = -\mathbf{F}(\mathbf{r}, t) , \quad (33)$$

$$\mathcal{D} [\mathbf{F}(\mathbf{r}, t)] = i\mathbf{F}(\mathbf{r}, t) . \quad (34)$$

These transformations for the operator fields are essentially the same as the transformation for the classical fields except for the time inversion shown in Eqs. (19) and (32). The reason for this difference is that in quantum mechanics, time inversion is represented in the Hilbert space by an *anti-unitary* operator that implies a complex conjugation (the necessity of this complex conjugation appears clearly upon observation of Schrödinger's or Dirac's equations). Therefore, the imaginary part of Eq. (15) changes sign twice; once due to the time inversion property of the magnetic field shown in Eq. (12) and a second time due to the required complex conjugation.

Let us postulate that the equal-time field commutators, that is, when both fields are evaluated at the same time t (not shown) but at different locations, have the form

$$[F_k^\dagger(\mathbf{r}') , F_l(\mathbf{r})] = \alpha_{kl}(\mathbf{r}' - \mathbf{r}) , \quad (35)$$

$$[F_k(\mathbf{r}') , F_l(\mathbf{r})] = \beta_{kl}(\mathbf{r}' - \mathbf{r}) , \quad (36)$$

where $\alpha_{kl}(\mathbf{r}' - \mathbf{r})$ and $\beta_{kl}(\mathbf{r}' - \mathbf{r})$ are two *functions*, not operators, that we must determine (of course at the right hand side of these equations an identity operator is implicit). In this *ansatz* we have imposed the condition that physical space is homogeneous because it doesn't matter at what space points the commutators are considered, that is, they can only depend on the separation $\boldsymbol{\rho} = \mathbf{r}' - \mathbf{r}$. The invariance under $\mathcal{C}$ transformation excludes the possibility that the commutator results in any odd power of the fields because the commutator involves the product of two fields and remains invariant under $\mathcal{C}$; therefore a term linear in the fields at the right hand side of Eq. (35) and (36) above is excluded (but not a bilinear form). If we perform the hermitian conjugation of Eq. (35) (that is, complex conjugation on the right hand side), with the usual rules for the conjugation of a product, we reach the conclusion that

$$\alpha_{lk}(-\boldsymbol{\rho}) = \alpha_{kl}^*(\boldsymbol{\rho}) . \quad (37)$$

Hermitian conjugation of Eq. (36) does not results in any condition for $\beta_{kl}(\boldsymbol{\rho})$.

Let us first require invariance of the commutation relation in Eq. (36) under the dual transformation. The right hand side is of course invariant but the left hand side changes

sign. Therefore we must have $\beta_{kl}(\boldsymbol{\rho}) = -\beta_{kl}(\boldsymbol{\rho})$, that is, $\beta_{kl}(\boldsymbol{\rho}) = 0$. This dual transformation places no constraints on the function $\alpha_{kl}(\boldsymbol{\rho})$ that will be fixed by other symmetry requirements. Let us now impose invariance under time inversion in Eq.(35) recalling that this transformation implies a complex conjugation. Eq.(32) indicates that the commutator remains unchanged and therefore we must have $\alpha_{kl}(\boldsymbol{\rho}) = \alpha_{kl}^*(\boldsymbol{\rho})$, that is, $\alpha_{kl}(\boldsymbol{\rho})$ is a *real* function and with this Eq.(37) becomes

$$\alpha_{kl}(\boldsymbol{\rho}) = \alpha_{lk}(-\boldsymbol{\rho}) . \quad (38)$$

Let us now look what the invariance under $\mathcal{P}$ implies. Let π denote the intrinsic parity of $\alpha_{kl}(\boldsymbol{\rho})$ under this transformation, that is, it may be a tensor or a pseudo tensor (it could even have mixed parity). Then, using Eq.(31) in the commutator of Eq.(35) we obtain

$$\alpha_{kl}(\boldsymbol{\rho}) = -\pi \alpha_{lk}(-\boldsymbol{\rho}) , \quad (39)$$

and therefore, comparing with Eq.(38) we conclude that $\pi = -1$, that is, $\alpha_{kl}(\boldsymbol{\rho})$ is a pseudotensor.

As a function of $\boldsymbol{\rho}$, $\alpha_{kl}(\boldsymbol{\rho})$ can be decomposed in an even (*Gerade*) part plus an odd (*Ungerade*) part under the change $\boldsymbol{\rho} \rightarrow -\boldsymbol{\rho}$, symbolically $\alpha = \alpha^G + \alpha^U$, and as a tensor, it can be decomposed in a symmetric part plus an anti-symmetric part under the permutation of the indices $kl \rightarrow lk$, symbolically $\alpha = \alpha^S + \alpha^A$. Therefore we have $\alpha = \alpha^{SG} + \alpha^{AG} + \alpha^{SU} + \alpha^{AU}$ but Eq.(38) implies that the *AG* and the *SU* part must vanish and we are left with only two parts $\alpha = \alpha^{SG} + \alpha^{AU}$.

Let us first analyse the *AU* part and we will later see that the *SG* part must vanish. It is easy to prove that any anti-symmetric tensor can be build by the contraction of ε_{kls} with an appropriately chosen vector. Therefore we have

$$\alpha_{kl}^{AU}(\boldsymbol{\rho}) = \varepsilon_{kls} \alpha_s(\boldsymbol{\rho}) , \quad (40)$$

where α_s is a component of an odd vector: $\alpha_s(\boldsymbol{\rho}) = -\alpha_s(-\boldsymbol{\rho})$ that we must still determine (we use the same letter α to denote the tensor and the vector but this should cause no confusion). Note that ε_{kls} is a pseudotensor and therefore $\alpha_{kl}^{AU}(\boldsymbol{\rho})$ has the required parity $\pi = -1$.

Let us now look at the symmetric-even part α^{SG} . As a matrix, it has the form

$$\alpha^{SG}(\boldsymbol{\rho}) = \begin{pmatrix} b_1 & a_3 & a_2 \\ a_3 & b_2 & a_1 \\ a_2 & a_1 & b_3 \end{pmatrix}$$

where a_i and b_i are *even* functions, that is, $a_i(-\boldsymbol{\rho}) = a_i(\boldsymbol{\rho})$ and $b_i(-\boldsymbol{\rho}) = b_i(\boldsymbol{\rho})$; but, if they don't vanish, we have a contradiction with the requirement that $\alpha_{kl}(\boldsymbol{\rho})$ is a pseudo tensor ($\pi = -1$), that is, it must change sign under a space inversion transformation $\mathcal{P}$. Therefore we must have $a_i = b_i = 0$. Summarizing, we have found that the commutation relations of the electromagnetic fields have the form

$$[F_k(\mathbf{r}') , F_l(\mathbf{r})] = [F_k^\dagger(\mathbf{r}') , F_l^\dagger(\mathbf{r})] = 0 , \quad (41)$$

$$[F_k^\dagger(\mathbf{r}') , F_l(\mathbf{r})] = \varepsilon_{kls} \alpha_s(\mathbf{r}' - \mathbf{r}) , \quad (42)$$

where $\alpha_s(\boldsymbol{\rho})$ is a component of an odd vector $\alpha_s(\boldsymbol{\rho}) = -\alpha_s(-\boldsymbol{\rho})$. In order to determine this vector let us use Eq.(27) and calculate a derivative ∂_s of any field component F_l with the condition that $s \neq l$, that is, $\delta_{sl} = 0$.

$$\begin{aligned} \partial_s F_l(\mathbf{r}) &= \frac{1}{i\hbar} [P_s , F_l(\mathbf{r})] \\ &= \frac{1}{i\hbar} \frac{-i}{16\pi c} \int d^3\mathbf{r}' \varepsilon_{skj} [F_k^\dagger(\mathbf{r}') F_j(\mathbf{r}') + F_j(\mathbf{r}') F_k^\dagger(\mathbf{r}') , F_l(\mathbf{r})] \\ &= \frac{-1}{16\pi\hbar c} \int d^3\mathbf{r}' \varepsilon_{skj} 2 F_j(\mathbf{r}') [F_k^\dagger(\mathbf{r}') , F_l(\mathbf{r})] \\ &= \frac{-1}{8\pi\hbar c} \int d^3\mathbf{r}' \varepsilon_{skj} \varepsilon_{klu} \alpha_u(\mathbf{r}' - \mathbf{r}) F_j(\mathbf{r}') \\ &= \frac{-1}{8\pi\hbar c} \int d^3\mathbf{r}' (-\delta_{sl}\delta_{ju} + \delta_{su}\delta_{jl}) \alpha_u(\mathbf{r}' - \mathbf{r}) F_j(\mathbf{r}') \\ &= \frac{-1}{8\pi\hbar c} \int d^3\mathbf{r}' \alpha_s(\mathbf{r}' - \mathbf{r}) F_l(\mathbf{r}') . \end{aligned}$$

This last line reminds us the definition of the derivative of Dirac's delta distribution. We have then

$$\alpha_s(\mathbf{r}' - \mathbf{r}) = 8\pi\hbar c \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) , \quad (43)$$

where the primed index s' indicates that we are deriving with respect to the components of $\mathbf{r}'$. With this we can finally write the equal time field commutators

$$[F_k(\mathbf{r}', t) , F_l(\mathbf{r}, t)] = [F_k^\dagger(\mathbf{r}', t) , F_l^\dagger(\mathbf{r}, t)] = 0 , \quad (44)$$

$$[F_k^\dagger(\mathbf{r}', t) , F_l(\mathbf{r}, t)] = 8\pi\hbar c \varepsilon_{kls'} \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) , \quad (45)$$

that written in terms of the more familiar electric and magnetic fields are,

$$[E_k(\mathbf{r}', t) , E_l(\mathbf{r}, t)] = [B_k(\mathbf{r}', t) , B_l(\mathbf{r}, t)] = 0 , \quad (46)$$

$$[E_k(\mathbf{r}', t) , B_l(\mathbf{r}, t)] = -i4\pi\hbar c \varepsilon_{kls'} \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) . \quad (47)$$

Now we will see that the commutation relations found are compatible with Maxwell's equations. For this, we can first calculate a simple expression for the tensor $M_{kls}(\mathbf{r}, t)$ in Eq.(30) using the commutators given in Eqs. (44) and (45).

$$M_{kls}(\mathbf{r}, t) = 16\pi\hbar c \varepsilon_{ksu'} \int d^3\mathbf{r}' \partial_{u'} \delta^3(\mathbf{r}' - \mathbf{r}) F_l(\mathbf{r}', t) \quad (48)$$

$$= -16\pi\hbar c \varepsilon_{ksu} \partial_u F_l(\mathbf{r}, t) . \quad (49)$$

Replacing it in Eq. (28) we find that the first Maxwell equation is identically satisfied whereas the second Maxwell Eq. (29) is satisfied, provided that $\partial_u F_u(\mathbf{r}, t) = 0$. This is a known fact, that the second Maxwell equation in quantum field theory must be introduced as a subsidiary condition. We can prove that this subsidiary condition is *necessary* for logical consistency, showing that otherwise we arrive at an absurd result:

$$\begin{aligned} \partial_u F_u(\mathbf{r}) &= \frac{1}{i\hbar} [P_u, F_u(\mathbf{r})] \\ &= \frac{1}{i\hbar} \frac{-i\varepsilon_{ukl}}{16\pi c} \int d^3\mathbf{r}' [F_k^\dagger(\mathbf{r}') F_l(\mathbf{r}') + F_l(\mathbf{r}') F_k^\dagger(\mathbf{r}'), F_u(\mathbf{r})] \\ &= \frac{-\varepsilon_{ukl}}{16\pi\hbar c} \int d^3\mathbf{r}' 2F_l(\mathbf{r}') [F_k^\dagger(\mathbf{r}'), F_u(\mathbf{r})] \\ &= \frac{-\varepsilon_{ukl}}{8\pi\hbar c} \int d^3\mathbf{r}' F_l(\mathbf{r}') 8\pi\hbar c \varepsilon_{kus'} \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) \\ &= (\delta_{uu} \delta_{ls'} - \delta_{us'} \delta_{lu}) \int d^3\mathbf{r}' F_l(\mathbf{r}') \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) \\ &= 2\delta_{ls'} \int d^3\mathbf{r}' F_l(\mathbf{r}') \partial_{s'} \delta^3(\mathbf{r}' - \mathbf{r}) \\ &= 2 \partial_l F_l(\mathbf{r}) . \end{aligned}$$

One can also show, after some lengthy calculations, that the commutation relations for the fields given in Eqs. (44) and (45) lead to the expected commutation relations for the hamiltonian and momentum, namely $[H, P_k] = [P_k, P_l] = 0$. In order to prove this we must use the subsidiary condition $\partial_u F_u(\mathbf{r}, t) = 0$ and that the rand terms in partial integrations vanish because the fields vanish at infinity.

With the knowledge of the equal time commutators given in Eqs. (44) and (45), we could calculate the commutators for different times performing a time evolution in one of the fields with the hamiltonian given in Eq.(24). For instance we have

$$[F_k^\dagger(\mathbf{r}', t'), F_l(\mathbf{r}, t)] = \left[\exp\left(\frac{i}{\hbar}(t' - t)H\right) F_k^\dagger(\mathbf{r}', t) \exp\left(\frac{-i}{\hbar}(t' - t)H\right), F_l(\mathbf{r}, t) \right] . \quad (50)$$

This would result in

$$[F_k(\mathbf{r}', t'), F_l(\mathbf{r}, t)] = -8\pi\hbar \varepsilon_{kls'} \partial_{s'} \partial_{t'} D(\mathbf{r}' - \mathbf{r}, t' - t) , \quad (51)$$

$$[F_k^\dagger(\mathbf{r}', t') , F_l^\dagger(\mathbf{r}, t)] = 8\pi\hbar \varepsilon_{kls'} \partial_{s'} \partial_{t'} D(\mathbf{r}' - \mathbf{r}, t' - t) , \quad (52)$$

$$[F_k^\dagger(\mathbf{r}', t') , F_l(\mathbf{r}, t)] = 8\pi\hbar c \left(\partial_{k'} \partial_{l'} - \frac{\delta_{kl}}{c^2} \partial_{t'}^2 \right) D(\mathbf{r}' - \mathbf{r}, t' - t) , \quad (53)$$

where

$$D(\boldsymbol{\rho}, \tau) = \frac{1}{4\pi\rho} (\delta(\rho + c\tau) - \delta(\rho - c\tau)) , \quad \rho = |\boldsymbol{\rho}| . \quad (54)$$

This is not a simple calculation and we will not perform it here. We just quote the result⁶ which, given in terms of the more familiar electric and magnetic fields, is

$$[E_k(\mathbf{r}', t') , E_l(\mathbf{r}, t)] = i4\pi\hbar c \left(\partial_{k'} \partial_{l'} - \frac{\delta_{kl}}{c^2} \partial_{t'}^2 \right) D(\mathbf{r}' - \mathbf{r}, t' - t) , \quad (55)$$

$$[B_k(\mathbf{r}', t') , B_l(\mathbf{r}, t)] = i4\pi\hbar c \left(\partial_{k'} \partial_{l'} - \frac{\delta_{kl}}{c^2} \partial_{t'}^2 \right) D(\mathbf{r}' - \mathbf{r}, t' - t) , \quad (56)$$

$$[E_k(\mathbf{r}', t') , B_l(\mathbf{r}, t)] = i4\pi\hbar \varepsilon_{kls'} \partial_{s'} \partial_{t'} D(\mathbf{r}' - \mathbf{r}, t' - t) . \quad (57)$$

These commutation relations were first obtained in 1928 by Jordan and Pauli⁷ but in the context of lagrangian quantum field theory.

The next step in the program of the quantization of the electromagnetic field would be to determine the eigenvalues and eigenvectors for the field operators. However, in the following section we will see that this is not meaningful and we will instead define another program consisting in the construction of the electromagnetic fields as emergent properties of a set of more fundamental entities.

V. INTERPRETATION AND CONCLUSION

In this section we will consider the physical meaning of the commutation relations of the electromagnetic field. Through the uncertainty principle, the commutation relations characterize the compatibility of the corresponding observables. That is, if the observables commute then there are quantum states where both observables are precisely determined; otherwise, the product of their indeterminacies has a lower bound proportional to the value of the commutator.

The function $D(\mathbf{r}' - \mathbf{r}, t' - t)$ defined in Eq.(54) vanishes everywhere except at the light cone, that is, except for those space time points $(\mathbf{r}', t')$ and $(\mathbf{r}, t)$ that can be connected by a light ray. Therefore the electromagnetic field for pairs of points that can not be joined by a light ray can assume exact values simultaneously because their commutator vanish. For points on the light cone, the commutator is highly singular; it involves second order derivatives of the singular distribution $D(\mathbf{r}' - \mathbf{r}, t' - t)$.

If these non vanishing commutators would take some *finite* value we would have a situation not unusual in quantum mechanics where the indeterminacy principle limits the simultaneous measurability of two observables but at least one of them could be known exactly and in all other cases both observables could be determined within some finite range of indeterminacy. Furthermore in the limit $\hbar \rightarrow 0$ we would recover the classical behaviour.

In the case of the electromagnetic fields, the *singular* character of the commutators have unacceptable consequences: assume that we could know the fields at one point $(\mathbf{r}, t)$; then they would propagate, even a short distance, to $(\mathbf{r}', t')$ where they would have an infinite indeterminacy. In this sense, the quantum mechanical treatment of the electromagnetic fields does not appear to be very reasonable. L. D. Landau and R. Peierls⁸ in 1931 recognized this difficulty and, without reference to the commutators but with some heuristic (but correct) arguments, reached the conclusion that for quantum theory “the field strength are not measurable quantities”. In an answer to this important remark, published two years later, N. Bohr and L. Rosenfeld⁹ noted that the commutation relations among *field averages*, taken on a finite volume, have non singular and finite values (depending on the size and shapes of the volumes where the averages are taken) and since every laboratory measurement of the fields involve an extended test charge, implying that what we really measure are field averages and not the fields at a point, they concluded that the difficulty can be dismissed because it has no consequence in a laboratory measurement. Rosenfeld commented¹⁰ that for them the “field components taken at a definite space time point are used in the formalism as an idealization without immediate physical meaning; the only meaningful statements of the theory concern averages of such field components over finite space time regions”.

The typically positivistic solution given by Bohr and Rosenfeld to the problem posed by the singular commutation relations of the fields is not satisfactory for those of us that believe that physics is not *only* concerned with real laboratory measurements but it also deals with idealizations and models that we build in order to describe reality. For this group there are two possibilities open: either is quantum mechanics wrong when applied to the electromagnetic field or the electromagnetic field is an essentially macroscopic system such that a quantum treatment of it is meaningless. Of course the enormous success of quantum mechanics in all realms of physics makes the second possibility more plausible. There are several examples of emergent properties of physical systems that do not make any sense at a deep microscopic level and whose quantization would be meaningless. For instance a temperature field $T(\mathbf{r}, t)$, or pressure or specific volume of a gas correspond to collective observables of large number of molecules. The molecules can, or must, be treated with quantum mechanics but to quantize pressure or temperature is meaningless. Already

at the classical level there is an indication that the electromagnetic field is not a basic or elementary system. This comes from the fact that the energy of the system does not scales with size of the system. Consider a physical system with fields $\mathbf{E}_1$ and $\mathbf{B}_1$ with an energy H_1 and another system with fields $\mathbf{E}_2$ and $\mathbf{B}_2$ with an energy H_2 . Now, the *compound* system with fields $\mathbf{E} = \mathbf{E}_1 + \mathbf{E}_2$ and $\mathbf{B} = \mathbf{B}_1 + \mathbf{B}_2$ will *not* have the energy $H_1 + H_2$ as one would expect from the combination of two elementary systems. The question is then, what are the corresponding “molecules” of the electromagnetic fields? The answer is well known: the photons.

Consequently we postulate that the electromagnetic field is a collective emergent properties of an ensemble of more fundamental entities, the photons, carriers of energy, momentum and angular momentum. The quantum mechanical description of the photons and how this set of bosons build the electromagnetic field will be the subject of following works.

It is relevant to mention that the choice adopted here, that the photons are the *primary ontology* that must be treated quantum mechanically and that the electromagnetic fields are macroscopic manifestations of an ensemble of them, is not a unique choice. There are indeed authors that consider the electromagnetic fields as the primary ontology and that the photons do not have an objective existence but are rather mathematical entities, “particle-like excitations”¹¹, corresponding to the normal mode decomposition of the fields. This alternative option can be adopted because, although the objective existence of the photons provide the most natural explanation of the photoelectric effect and of Compton scattering, these are not compelling evidence for their existence because these effects can also be explained by a semiclassical theory combining unquantized fields with quantum theory of matter^{12,13}. The existence of these two alternative interpretations indicate that the quantum description of electromagnetic radiation is not a closed subject and that Einstein’s question *What are light quanta?* has not been answered.

ACKNOWLEDGMENTS

I would like to thank H. Martín for challenging discussions and E. Fernández for bibliographic assistance. This work received partial support from “Consejo Nacional de Investigaciones Científicas y Técnicas” (CONICET).

REFERENCES

- ¹ “Heute glaubt zwar jeder Lump, er wisse es, aber er täuscht sich” in *Albert Einstein-Michele Besso Correspondence 1903-1955* P. Speziali, Ed. Hermann, Paris, 1972. Letter dated 12 December 1951, Pg.453. The English translation of the German word “Lump” has a large dispersion. I have adopted one that is close to the French word “abrutí” chosen by the editor P. Speziali.
- ² For instance L. I. Schiff *Quantum Mechanics* Mc Graw-Hill Kogakusha, Tokio, 1968.
- ³ The Heaviside system is the “right” one where the electric and magnetic fields have the same units, as they should because they are different components of the same tensor, and the 4π is in Coulomb law, where it belongs because of its inherent spherical symmetry. Unfortunately, the SI system of units has hit the wrong choice.
- ⁴ See D. Hertens “Oersted Medal Lecture 2002: Reforming the mathematical language of physics” *Am. J. Phys.* **71**, 104-121, (2003), and references therein.
- ⁵ See for instance Eq.(7.3.28) and (7.3.29) in S. Weinberg *The Quantum Theory of Fields I* Cambridge U. Press, Cambridge, 1995
- ⁶ See for instance reference 2 above or F. J. Ynduráin, *Relativistic Quantum Mechanics and Introduction to Field Theory* Springer Verlag, Berlin, 1996.
- ⁷ P. Jordan and W. Pauli “Zur Quantenelectrodynamik ladungsfreier Felder” *Z. Physik* **47**,151-173 (1928)
- ⁸ L. D. Landau and R. Peierls, “Erweiterung des Unbestimmtheitsprinzips für die relativistische Quantentheorie”, *Z. Phys.* **69**, 56 (1931). A translation of this work can be found in *Quantum Theory and Measurement* J. A. Wheeler and W.H. Zurek Eds. Princeton University Press, Princeton, 1983.
- ⁹ N. Bohr and L. Rosenfeld, “Zur Frage der Messbarkeit der electromagnetischen Feldgrößen”. *Mat.-fys. Medd. Dan. Vid. Selsk.*, 12, 8 (1933). A translation of this work can be found in *Quantum Theory and Measurement* J. A. Wheeler and W.H. Zurek Eds. Princeton University Press, Princeton, 1983.
- ¹⁰ L. Rosenfeld, “The Measurability of the Electromagnetic Field”, in *Quantum Theory and Measurement* J. A. Wheeler and W.H. Zurek Eds. Princeton University Press, Princeton, 1983.

- ¹¹ L.E. Ballentine, *Quantum Mechanics* World Scientific, Singapore 1998, pages 526, 540. See also G. Greenstein and A.G. Zajonc *The Quantum Challenge* Jones and Bartlett Pub., Sudbury MA. 1997, page 35.
- ¹² M. O. Scully and M. Sargent III “The concept of the photon” *Physics Today*/March 1972, 38-47, 1972.
- ¹³ R. Kidd, J. Ardin and A. Anton “Evolution of the modern photon” *Am. J. Phys.* **57** 27-35, 1989.